

# Synthesis and Characterization of Novel N-Substituted-3-chloro-2-Azetidinones as Potential Anticonvulsant Agents.

Vijay Kumar, M.M.J.,\*<sup>I</sup> Yogananda, R., <sup>II</sup> Snehalatha., <sup>II</sup> Shameer, H., <sup>III</sup> Jayachandran, E., <sup>III</sup> Sreenivasa, G.M. \*\*<sup>III</sup>

<sup>1</sup>Dept of Pharm. Chemistry, <sup>11</sup>Dept of Pharm. Ceutics, S.J.M. College of Pharmacy, Chitradurga-577502, Karnataka, India. <sup>111</sup>P.G. Dept. of Pharm. Chemistry, S.C.S. College of Pharmacy, Harapanahalli–583131, Karnataka, India.

#### ABSTRACT

Various substituted 4(m-hydroxy-p-methoxy phenyl)-1[(6'-fluoro-7'-substituted (1,3)-benzothiazol-2'-yl) amido-2phenyl] 3-chloro azetidin–2–one containing different functional groups have been synthesized. The lead compounds were characterized by melting point, TLC, calculated elemental analysis, UV, IR, <sup>1</sup>HNMR and Mass spectral studies. The compounds were tested for Anticonvulsant studies by PTZ induced method and showed significant activity at low and high concentration compared to standard; still further studies are requested.

Keywords: Anticonvulsant activity, Azetidinone, Benzothiazole, Fluorine.

#### **INTRODUCTION**

2-Azetidinones, commonly known as βlactams. are well-known heterocyclic organic and compounds among the medicinal chemists.<sup>1</sup> The activity of the famous antibiotics such as penicillins, cephalosporins and carbapenems are attributed to the presence of 2-azetidinone ring in them. Recently, some other types of biological activity besides the antibacterial activity have been reported in compounds containing 2-azetidinone ring.<sup>2,3</sup> Such biological activities include antimicrobial,<sup>4</sup> anti-tuburcular,<sup>5</sup> carbonic anhydrase inhibitors,<sup>6</sup> local anaesthatics,<sup>7</sup>

anti-inflammatory,<sup>8</sup> anthelmintic,<sup>9</sup> anticonvulsant,<sup>10</sup> hypoglycemic agents activity.<sup>11</sup> The  $\beta$ -lactams also serve as synthons for many biologically important classes of organic compounds.<sup>12</sup> Due to this, the investigation of chemistry and biology of these compounds continue to appeal the synthetic and medicinal organic chemists.<sup>1-4</sup> It is well known that the introduction of fluorine atom into an organic molecule causes dramatic changes in its biological profile, mainly due to high electro negativity of fluorine, the strong carbon-fluorine bond and increased solubility in lipids.<sup>13</sup> Therefore it was thought worthwhile to synthesize better kinds of drugs by incorporating azetidinone in benzothiazole moiety.

In search for new biodynamic potent molecule, it was thought worthwhile to incorporate some additional heterocyclic moieties in the  $\beta$ -lactam nucleus and study their biological and pharmacological activity.<sup>14</sup> The review of literature reveal prompted us to synthesize substituted fluorobenzothiazole, azetidinone targeted compounds and those will be screened for anticonvulsant activity.

### MATERIALS AND METHODS Chemicals and Reagents

4-fluoro-3-chloro aniline, Potassium thiocyanate, Glacial acetic acid, Bromine, Anthranillic acid, Pyridine, Vanillin, Ethanol, Conc. Hydrochloric acid, Chloroacetyl chloride, Triethylamine,

For Correspondence:

<sup>\*</sup>E-mail: vijaykumarmmj@yahoo.in

<sup>\*\*</sup>E-mail : gms 2006@rediffmail.com

N,N<sup>\dimethyl</sup> formamide (DMF), various substituted aniline, morpholine, piperazine and diphenylamine.

# **Experimental Section**

**Step I:** 4-fluoro-3-chloro aniline was treated with potassium thiocyanate (KSCN) in presence of glacial acetic acid and bromine to get 2-amino-6-fluoro-7-chlorobenzothiazole.

**Step II:** 2-amino-6-fluoro-7-chlorobenzothiazole treated with Anthranillic acid in presence of Pyridine to get 2 (o-amino phenyl amido) 6–fluoro -7-chloro (1,3) benzothiazole.

**Step III:** 2 (o-amino phenyl amido) 6– fluoro -7-chloro (1,3) benzothiazole reflexed with vanillin and alcohol in presence of Conc.HCl to get 2 (3-hydroxy-4-methoxy benzylidene amino phenyl amido) 6-fluoro-7-chloro-(1,3) benzothiazole or Schiff's base.

**Step IV:** A Solution of Schiff's base (0.01 mol) in 1,4-dioxane (50ml) was added to well-stirred mixture of Chloroacetyl Chloride (0.95 ml, 0.012 mol) and Triethylamine (1.08 ml, 0.02 mol) at 0° C. The reaction mixture was then stirred for 18 - 20 hrs and kept aside for 3 days at room temperature. The product was recrystallised from N,N' Dimethyl formamide (DMF).

**Step V:** Azetidine were treated with double the quantities of various substituted aniline, anisidine, PABA and piperazine, refluxed for 2 hours in presence of N,No-dimethyl formamide (DMF). The mixture was cooled and poured in to crushed ice. The solid separated was filtered off, dried and crystallized from alcohol and benzene.

## **General Procedures**

Melting points were determined in open capillaries and are uncorrected. IR spectra (KBr pellet technique) were recorded using a Perkin-Elmer 237 spectrophotometer. <sup>1</sup>H NMR spectra were recorded on Bruker AM 400 instrument (at 400 MHz) using tetramethylsilane (TMS) as an internal

standard and DMSO-d6 as a solvent. Chemical shifts are given in parts per Splitting patterns million (ppm). are designated as follows: s- singlet, d- doublet, t- triplet, q- quartet and m-multiplet. Mass spectra (MS) were recorded on Schimadzu GC-MS operating at 70eV. All the synthesized compounds were purified by recrystallization. The reactions were followed up and the purity of compounds was monitored on pre-coated TLC plates and visualizing the spots in ultraviolet light.

## In Vitro Anticonvulsant Study

In the present study the mice of either sex, weighing between 18-25 g were selected and divided into control, test and standard.

Before experiment the animal were fasted for 24 hrs with only water *ad-libitum*. Control group received only 0.5 ml DMF as vehicle. Standard group animals were received diazepam (4 mg/kg b.w.) oral test group animals were received the synthesized derivatives at 4 mg/kg b.w. oral in DMF.

Now for the animals of control group pentylene tetrazole (PTZ) 1 ml / 100 g b.w. was administered and actions like straubs tail, jerky movements of whole body and conclusions were observed.

For animals of standard test group PTZ was injected (1 ml/100 g b.w.). After 30 mins animals of standard and test received diazepam and synthesized derivatives respectively.<sup>15</sup> Observations were made and results were tabulated.

## **RESULTS AND DISCUSSION**

Synthesis and pharmacological screening of 4(m-hydroxy-p-methoxy phenyl)-1[(6'-fluoro-7'-substituted (1,3)-benzothiazol-2'-yl) amido-2-phenyl] 3-chloro azetidin–2–one were tested for anticonvulsant activity by PTZ induced method compared to standard Diazepam; showed significant anticonvulsant activity.

Among compounds tested  $A_{3}$ ,  $A_{7}$ ,  $A_{8}$  and  $A_{12}$  showed significant anticonvulsant activity.

### Scheme



# Table No. 1 Anti Convulsant Activity (PTZ – Induced)

Stock Solution. Test Drugs 5 mg/ml Mice (Body weight- 18-25gm) Dose: PTZ – 80 mg/kg I.P.

Animal – Albino PTZ – 8 mg/ml Test Drugs – 50 mg / kg orally

CI		Dece		Death /		
No	Treatment	(ma/ka)	Onset	Nature of	Clonic time	
NO.		(iiig/kg)	(Sec)	Severity	in Sec	Recovery
1			43		55	
2	Water		75	lorky.	103	
3	+	5 ml / kg + 80	63	Movement	95	5/5
4	PTZ		55	MOVEMENT	75	
5			45		55	
1			67		85	
2	A 1		71	lorlar	93	
3	+	50 + 80	55	Jerky	95	5/5
4	PTZ		56	wovernent	75	
5			58		95	
1			71		85	
2	A 2		62	le ulu i	83	
3	+	50 + 80	52	Јегку	95	5/5
4	PTZ		63	wovement	85	
5			45		85	
1			93		450	
2	A 3		102	Straub's Tail	451	0/5
3	+ PTZ	50 + 80	103		220	
4			83		395	
5			95		423	
1			93	Jerky Movement	85	5/5
2	Ae	50 + 80	102		73	
3	+		103		95	
4	PTZ		83		95	
5			95		75	
1			63		85	
2	Α-7		52	1	93	
3		50 + 80	55	Jerky	95	3/5
4	PTZ		71	Movement	105	
5			63		55	
1			83		450	
2	A。		93		451	
3	+	50 + 80	105	Straub's Tail	220	4/5
4	PTZ	00100	71		390	1,0
5			103	1	423	
1			53		85	
2	A 10		55	1.	73	
3	+	50 + 80	55	Jerky	95	5/5
4	PT7		79	Movement	75	
5			65	1	95	

1			65		95	
2	A 11		63	lorlar	83	5/5
3	+	50 + 80	67	Movement	85	
4	PTZ		73		75	
5			81		95	
1			52		85	
2	A <sub>12</sub> + PTZ		53	lorla	83	4/5
3		50 + 80	55	Movement	95	
4			65		95	
5			75		75	
1			97		445	
2	Diazapam + PTZ		105		455	
3		+ 50 + 80 PTZ	110	Straub's Tail	210	0/5
4			85		385	
5			93		415	

 Table No. 2
 Analytical Data

SI.	Compound	<b>M.P</b> /	%	MOL FORM	M Wt	Calculated %			
No	Code	B.P°C	Yield	MOL. FORM	1 <b>11. VV L.</b>	С	Н	Ν	
1	$A_1$	190	78%	$C_{30}H_{21}O_6SN_5FCl$	634	56.83	3.34	11.05	
2	A <sub>2</sub>	178	82%	$C_{30}H_{21}O_6SN_5FCl$	634	56.83	3.34	11.05	
3	A <sub>3</sub>	183	75%	$C_{30}H_{21}O_6SN_5FCl$	634	56.83	3.34	11.05	
4	$A_4$	164	72%	$C_{30}H_{21}O_4SN_4FCl_2$	623	57.79	3.39	8.99	
5	A <sub>5</sub>	132	74%	$C_{30}H_{21}O_4SN_4FCl_2$	623	57.79	3.39	8.99	
6	$A_6$	126	73%	$C_{30}H_{21}O_4SN_4FCl_2$	623	57.79	3.39	8.99	
7	$A_7$	112	76%	$C_{30}H_{22}O_4SN_4FCl$	589	61.17	3.76	9.51	
8	$A_8$	124	65%	$C_{31}H_{24}O_5SN_4FCl$	619	60.14	3.91	9.05	
9	A <sub>9</sub>	118	69%	$\mathrm{C_{31}H_{24}O_5SN_4FCl}$	619	60.14	3.91	9.05	
10	$A_{10}$	158	83%	$C_{31}H_{24}O_5SN_4FCl$	619	60.14	3.91	9.05	
11	A <sub>11</sub>	260	77%	$C_{31}H_{22}O_6SN_4FCl$	633	58.82	3.50	8.85	
12	A <sub>12</sub>	308	85%	$C_{28}H_{25}O_4SN_5FCl$	582	57.78	4.33	12.03	

Compound	Ar-NH (in cm <sup>-1</sup> )	C=0 Stretching (in cm <sup>-1</sup> )	C=N Stretching (in cm <sup>-1</sup> )	C=C Stretching (in cm <sup>-1</sup> )	NO <sub>2</sub> (in cm <sup>-1</sup> )	C-F (in cm <sup>-1</sup> )	C – S Stretching (in cm <sup>-1</sup> )	Sec.Ar. Amine (in cm <sup>-1</sup> )	C – Cl Stretching (in cm <sup>-1</sup> )	C-O-C Stretching (in cm <sup>-1</sup> )	Ar-OH Stretching (in cm <sup>-1</sup> )
$A_1$	3350	1750	1550	1710	1450	1130	720	1300	840	1250	1390
A <sub>2</sub>	3370	1710	1525	1680	1450	1160	720	1340	840	1250	1390
A <sub>3</sub>	3370	1700	1540	1660	1420	1160	725	1310	850	1255	1380
A <sub>4</sub>	3380	1730	1540	1680	-	1155	720	1300	850	1250	1380
A <sub>5</sub>	3400	1765	1540	1690	-	1170	725	1310	820	1250	1380
A <sub>6</sub>	3290	1720	1530	1680	-	1160	725	1300	840	1250	1380
A <sub>7</sub>	3390	1755	1510	1690	-	1150	720	1255	840	1220	1380
A <sub>8</sub>	3350	1720	1540	1685	-	1165	725	1310	830	1250	1390
A <sub>9</sub>	3310	1730	1550	1650	-	1130	725	1310	840	1245	1380
A <sub>10</sub>	3400	1750	1560	1660	-	1170	730	1300	850	1230	1385
A <sub>11</sub>	3320	1700	1530	1640	-	1165	730	1310	840	1270	1380

# Table No. 3 Characteristics IR absorption bands :

SI No.	Compound Code	Hydrogen	δ(ppm)	Multiplity	Solvent
1	$A_3$	-Ar-H- -NH- β lactum 2H – Proton	7.0 – 7.8 5.4 6.6	Multiplet Singlet Doublet	CDCl <sub>3</sub>
2	$A_6$	-Ar-H- - NH - β lactum 2H – Proton	7.0 – 7.8 5.3 6.6	Multiplet Singlet Doublet	CDCl <sub>3</sub>

Table No. 4 <sup>1</sup>H NMR Spectral Data



Fig 1. NMR Spectra - A<sub>6</sub>

Sl no	Compound Code	Calc. Mol. weight	Mol Formula	Fragmentation	m/z
1	A <sub>3</sub>	634.03	C <sub>30</sub> H <sub>21</sub> O <sub>6</sub> SN <sub>5</sub> FCl	$ \begin{array}{c} M^{+1}(CH_{3}O, Cl, NO_{2}) \\ M^{+2} (C_{6}H_{3}, C_{3}NO) \\ M^{+3} \{(C_{6}H_{4})_{2}O\} \end{array} $	517.5 379.4 201.3
2	$A_6$	623.48	C <sub>30</sub> H <sub>21</sub> O <sub>4</sub> SN <sub>4</sub> FCl <sub>2</sub>	$ \begin{array}{l} M^{+1}(\text{-CH}_3) \\ M^{+2} \{C_6H_3(\text{OH})\text{-O}, \text{Cl}, \text{O}\} \\ M^{+3} (\text{N-C-C-C}_9\text{Cl}) \\ M^{+4} \{(C_6H_4)_2\} \end{array} $	613.9 454.7 369.8 203.2

Table No. 5 Mass Spectral Data





### CONCLUSION

Result of present study demonstrate that, a new class of different aromatic aniline, anisidine, PABA, piperzine, encompassing azetidinone derivatives were synthesized and evaluated as antibacterial agents. The newly synthesized heterocyclics exhibited promising anticonvulsant activity using PTZ induced method. The anticonvulsant studies showed significant activity at low and high concentration compared to standard. It can be concluded that this class of compounds certainly holds great promise towards good active leads in medicinal chemistry. А further study to acquire more information concerning pharmacological activity is in progress.

## ACKNOWLEDGEMENT

The authors are thankful to Shri. Sha. Bra. Chandramouleshwara Shivacharya swamiji, President, T.M.A.E. Society, Harapanahalli, for providing necessary facilities through the Principal, Dr. S.Ramachandra Setty, S.C.S. College of Pharmacy, Harapanahalli to carry out this work.

### References

- [1] Singh, G. S., *Tetrahedron* 2003, 59, 7631.
- [2] Singh, G. S., *Mini-Rev. Med. Chem.* 2004, 4, 69.
- [3] Singh, G. S., *Mini-Rev. Med. Chem.* 2004, 4, 93.
- [4] Chalreck, P., Synthesis of 2-styryl benzothiazolinium salts for antibacterial and fungicidal activity; Chem. Abstr., 1988, 109, 3771 x.
- [5] Ares Jaffrey., Synthesis of 2-substituted benzothiazoles from 2-fluoro phenyl

*isothiocynate, Chem. Abstr.*, 1991, 114, 71453 d.

- [6] Wollesdrof, O.W. Jr., Schwam, H., Synthesis of 1-o-acyl derivatives of hydroxyl benzothiazol 2-sulfonamide as topically active carbonic-anhydrous inhibitors, Chem. Abstr., 1989, 111, 194656 x.
- [7] Costakes, E., Tsatsas. G., Synthesis of 2-(alkylamino acyl imino) 3-methyl benzothiazolines for local anaesthetic activity, Chem. Abstr., 1979, 90, 203935 q.
- [8] Sreenivasa Rao, D., Jayachandran, E., Sreenivasa, G.M., Shivakumar, B., Inhibition of albumin denaturation and anti-inflammatory activity of 2-[N-p-Tolyl sulphon hydrazino]-6-fluoro-7substituted (1,3) benzothiazoles, Oriental J. of Chemistry, 2005, 21(1), 113-116.
- [9] Sreenivasa, G.M., Shivkumar, B., Jayachandran. E., Anthelmintic activity of 8-fluoro-9-substituted (1,3)benzothiazolo (5,1-b)-1,2,4-triazole on perituma posthuma, Indian Drugs, 2006, 43 (4).
- [10] Abdel Ghany aly el-helby., Mohammed Hemeda abdel wahed., *Acta Pharm.*, 2003, 53, 127-138.
- [11] Chernykh, V.P., Sidorenko, O.F., Synthesis of ethyl N-[6-substituted benzo (tetrahydrobenzo) 2-thiazolyl] oxamate for hypoglycemic activity, Chem. Abstr., 1983, 98, 89233 x.
- [12] Conte, L., et al., Liquid-Phase fluorination of aromatic compounds by elemental fluorine, J. Fluorine Chem, 1995, 70, 175-179.

- [13] Desbois Michel., Eur Pat., APP.1 EP. 248746 (IPC Co7Co85/00), Synthesis of 4-fluoro-3-chloroaniline from 4-halo nitrobenzene, Chem. Abstr., 1987, 108, 159896 a.
- [14] Biological assays., Indian Pharmacopoeia Published by Govt. of India, 1996, 2, 88.
- [15] Turner, R.A., Anti-epileptic drugs, in screening methods in pharmacology, XV edn., (Ed. Turner RA), Academic Press, New York, 1965, 165-172.